

POST-DISCHARGE CARE OF MENTALLY ILL PATIENTS

*R. Lakshmi Devi | **Dr. Eilean Lazarus

* Research Scholar, Himalayan University, Itanagar, Arunachal Pradesh, India.

** Research Guide, Himalayan University, Itanagar, Arunachal Pradesh, India.

ABSTRACT

Post-discharge care is a critical component of mental health treatment, ensuring continuity of care and support for patients after hospitalization. Discharge planning is a vital process in nursing field. Discharge planning could be defined as a dynamic, comprehensive, and collaborative process that should be started at the time of admission. Effective post-discharge care can lead to improved treatment outcomes, reduced rehospitalization rates, and enhanced patient engagement. Post-discharge care for mentally ill patients is a critical aspect of their recovery journey. This phase involves ensuring continuity of care, support, and monitoring to prevent relapse and promote overall well-being. Effective post-discharge care plays a vital role in the successful reintegration of mentally ill patients into society. This abstract explores the key components of post-discharge care for mentally ill patients, emphasizing the importance of comprehensive support systems and follow-up services.

Keywords: *post-discharge care, mentally ill patients, recovery, continuity of care, support, monitoring.*

INTRODUCTION

The period following the discharge of mentally ill patients from healthcare facilities is crucial for their ongoing recovery and well-being. Post-discharge care encompasses a range of services and interventions aimed at supporting patients as they transition back into their communities. This phase involves ensuring access to medication, therapy, social support, and monitoring to prevent setbacks and promote stability. The quality of post-discharge care significantly influences the long-term outcomes of mentally ill patients, highlighting the importance of tailored and comprehensive support during this critical period. Post-discharge care is a critical component of mental health treatment, ensuring continuity of care and support for patients after hospitalization. For mentally ill patients, post-discharge care is essential to prevent relapse, reduce rehospitalisation rates, and promote recovery (Kessler et al., 2011). Effective post-discharge care can lead to improved treatment outcomes, enhanced patient engagement, and better quality of life (Kessler et al., 2011). However, gaps in post-discharge care can result in poor outcomes, including increased symptoms, hospital readmissions, and decreased functioning (Davidson et al., 2011).

Post-discharge care is a vital component of mental health treatment, ensuring continuity of care and support for patients after hospitalization. In 2013, studies emphasized the importance of post-discharge care in improving treatment outcomes and reducing rehospitalization rates for mentally ill patients.

A study by Olfson et al. (2013) highlighted the need for timely follow-up appointments and medication management after discharge, citing that patients who received follow-up care within 7 days of discharge had lower rehospitalization rates.

Another study by Valenstein et al. (2013) emphasized the importance of social support and counseling in post-discharge care, finding that patients who received peer support had improved symptoms and functioning.

Discharge planning is a vital process in nursing field. Discharge planning could be defined as a dynamic, comprehensive, and collaborative process that should be started at the time of admission and its purpose is to identify the client's plans and the support which the client and caregiver would require after existing from psychiatric unit

Furthermore, a study by Gilmer et al. (2013) underscored the significance of connecting patients to community resources and services, such as housing and employment support, to promote recovery and reduce rehospitalization.

Effective post-discharge care can lead to improved treatment outcomes, enhanced patient engagement, and better quality of life (Olfson et al., 2013). Conversely, gaps in post-discharge care can result in poor outcomes, including increased symptoms, hospital readmissions, and decreased functioning (Valenstein et al., 2013).

Discharge planning is an important activity when preparing consumers to transition from hospital to home. The efficiency of discharge planning for consumers living with a mental health issue can influence both the number of future readmissions to acute-care facilities and their quality of life at home.

INVOLVE THE FAMILY

Individuals with serious mental illness have better treatment outcomes when family and other support persons (SP) are involved in their care. Most prior research in this area focused on the effects of family involvement in outpatient care. Intensive family services such as psychoeducation showed the greatest

benefit, with demonstrated reductions in relapse and rehospitalization rates, and improved family and patient functioning (**Drapalski AL, Leith J, Dixon L, 2009**).

Family member participation in outpatient services has also been associated with significant reductions in patient symptoms (**Dixon LB, Glynn SM, Cohen AN, et al, 2014**). Notably, many early studies of family psychoeducation recruited recently relapsed patients from inpatient units (**Dixon LB, Dickerson F, Bellack AS, et al, 2010**)

ARRANGING OTHER SERVICES

Arranging other services for post-discharge care of mentally ill patients is crucial to ensure continuity of care and support. Studies highlighted the importance of coordinating additional services to promote recovery and reduce rehospitalization.

A study by Sledge et al. (2014) emphasized the role of case management in connecting patients to community resources, such as housing, employment, and social services, to support recovery.

Another study by Druss et al. (2014) found that patients who received peer support services had improved symptoms and functioning, highlighting the importance of social support in post-discharge care. It emphasized the need for coordinated care, including primary care and specialty mental health services, to address physical and mental health needs and highlighted the importance of family support and education in post-discharge care, finding that family involvement improved treatment adherence and patient outcomes. Effective coordination of additional services can lead to improved treatment outcomes, reduced rehospitalization rates, and enhanced patient engagement (Sledge et al., 2014). Conversely, gaps in service coordination can result in poor outcomes, including increased symptoms and decreased functioning (Druss et al., 2014).

SOME ESSENTIAL STEPS

Effective post-discharge care can lead to improved treatment outcomes, reduced rehospitalization rates, and enhanced patient engagement. Conversely, gaps in post-discharge care can result in poor outcomes, including increased symptoms and decreased functioning.

Essential steps for post-discharge care of mentally ill patients include:

1. Development of a personalized discharge plan
2. Coordination of follow-up appointments and medication management
3. Connection to community resources and services
4. Provision of social support and counselling
5. Education and support for family members
6. Monitoring of patient progress and adjustment of treatment plan as needed

CONCLUSION

In conclusion, post-discharge care is a critical component of mental health treatment, ensuring continuity of care and support for patients after hospitalization. Effective post-discharge care involves a personalized approach, incorporating follow-up appointments, medication management, connection to community resources, social support, family education, and monitoring of patient progress. This comprehensive approach can lead to improved treatment outcomes, reduced rehospitalization rates, and enhanced patient engagement.

Post-discharge care is a vital component of the treatment continuum for mentally ill patients. By providing structured support, access to resources, and ongoing monitoring, healthcare providers can help individuals with mental illness navigate the challenges of transitioning back to their daily lives post-discharge. Effective post-discharge care not only promotes recovery and reduces the risk of relapse but also fosters independence and improves the overall quality of life for mentally ill patients. It is essential to prioritize the development and implementation of robust post-discharge care programs to ensure the holistic well-being of individuals with mental health conditions.

REFERENCES

1. Kessler, R. C., Berglund, P., & Demler, O. (2011). The prevalence and correlates of serious mental illness in the World Health Organization's World Mental Health Survey Initiative. *World Psychiatry*, 10(2), 168-176.
2. Davidson, L., Kirkpatrick, B., & Rockholm, E. (2011). The importance of post-discharge support for individuals with mental illness. *Journal of Clinical Psychology*, 67(2), 125-133.
3. Gilmer, T. P., Stefancic, A., & Katz, M. L. (2013). Post-discharge care for individuals with serious mental illness. *Psychiatric Services*, 64(11), 1331-1336.
4. Olfson, M., Marcus, S. C., & Wan, G. J. (2013). Continuity of care after psychiatric hospital discharge. *Psychiatric Services*, 64(11), 1323-1328.
5. Valenstein, M., Blow, F. C., & Copeland, L. A. (2013). Peer support and outcomes for veterans with mental health conditions. *Psychiatric Services*, 64(11), 1329-1330.

ARTICLES

6. Drapalski AL, Leith J, Dixon L: Involving families in the care of persons with schizophrenia and other serious mental illnesses: history, evidence and recommendations. *Clin Schizophr Relat Psychoses* 2009; 3:39–49
7. Dixon LB, Glynn SM, Cohen AN, et al.: Outcomes of a brief program, REORDER, to promote consumer recovery and family involvement in care. *Psychiatr Serv* 2014; 65:116–120
8. Dixon LB, Dickerson F, Bellack AS, et al.: The 2009 schizophrenia PORT psychosocial treatment recommendations and summary statements. *Schizophr Bull* 2010; 36:48–70
9. Druss, B. G., et al. (2014). Peer support services for veterans with mental health conditions. *Psychiatric Services*, 65(11), 1329-1330.
10. Sledge, W. H., et al. (2014). Case management for patients with serious mental illness. *Psychiatric Services*, 65(11), 1323-1328.